

Fairfield County Implants & Periodontics, LLC

POST-OPERATIVE INSTRUCTIONS

MEDICATIONS	O Ibuprofen 600 mg every 3-4 hours for first 3 days <i>NO MATTER WHAT</i> . Next doses at: O Take extra pain med like Vicodin® (hydrocodone) <i>IN ADDITION TO</i> ibuprofen if needed Continue antibiotics until gone Continue Arnica until gone (5 pellets, 3x/day) IF GIVEN, apply Arnica cream externally on face 3x/day
SWELLING	Swelling is normal and peaks 48 to 72 hours post-procedure O Use ice pack (15 min on/15 min off) O Sleep on two pillows to elevate head
BLEEDING	Light bleeding may occur, even from the nose, up to 72 hours post-procedure O Control by applying pressure with moist gauze or wet tea bag for 15 min
NASAL CONGESTION	O <i>DO NOT</i> blow your nose O May use oral decongestant like Sudafed® O Avoid nasal spray
FOOD	O Eat soft foods such as: • Yogurt or cottage cheese • Soup • Well-cooked veggies • Soft bread • Mashed potatoes • Stuffing • Pudding or gelatin • Sorbet • Oatmeal • Porridge • Pasta • Eggs • Applesauce • Bananas • Protein shakes • Fish O DO NOT eat chewy, hard, fried or spicy food like cereal, crusty bread, pizza, candy, corn, rice, popcorn, nuts, gum, steak, wings, sausage, chips, nachos, pretzels, seeds
ORAL HYGIENE	O Rinse with Peridex® (0.12% chlorhexidine) or warm salt water 2x/day O Avoid brushing or flossing the surgical area O OK to clean other teeth
EXERCISE	Avoid for a minimum of 48 hours
DENTURES	Avoid wearing temporary denture as much as possible
DO NOT	O Smoke O Spit out with force O Use a straw O Remove sutures yourself O Remove dressing if applied

Please contact us anytime if you have any questions or concerns before or after your procedure. If you have any questions after hours, please call the office number (below) and follow the prompts. Thank you.

Dr. Michael Sonick (203) 254-2006 (office) mike@sonickdmd.com Dr. "Ray" Rui Ma (203) 254-2006 (office) ray@sonickdmd.com